



US006468542B2

(12) **United States Patent**
Liu et al.

(10) **Patent No.:** US 6,468,542 B2
(45) **Date of Patent:** Oct. 22, 2002

(54) **GERMINATION ACTIVATED *GANODERMA LUCIDUM* SPORES AND METHOD FOR PRODUCING THE SAME**

(76) **Inventors:** Xin Liu, Building No. 391, 135
Xingang Xi Road, Guangzhou (CN);
Chee-Keung Chung, Room 1505,
Argyle Centre, 688 Nathan Rd.,
Mongkok, Kowloon (HK)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** 09/802,862

(22) **Filed:** Mar. 12, 2001

(65) **Prior Publication Data**

US 2001/0031265 A1 Oct. 18, 2001

Related U.S. Application Data

(62) Division of application No. 09/524,508, filed on Mar. 13,
2000, now Pat. No. 6,316,002.

(60) Provisional application No. 60/158,377, filed on Oct. 12,
1999.

(51) **Int. Cl.⁷** A61K 35/84

(52) **U.S. Cl.** 424/195.15

(58) **Field of Search** 424/195.15

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,595,742 A * 1/1997 Fujiwara et al.

FOREIGN PATENT DOCUMENTS

CN	1101860	* 10/1993
CN	1111529	* 11/1995
CN	1118005	* 3/1996
CN	1122658	* 5/1996
CN	1134306	* 10/1996
CN	1165032	* 11/1997

* cited by examiner

Primary Examiner—Christopher R. Tate

(74) *Attorney, Agent, or Firm*—Fei-Fei Chao; Venable,
Baetjer, Howard, and Civiletti

(57) **ABSTRACT**

The present invention describes a method for germination activating spores of *Ganoderma lucidum* to produce bioactive substances which has medicinal effects on patients with immunological disorders, cancer, AIDS, hepatitis, diabetes, and cardiovascular diseases, and can prevent or inhibit free radical oxidation and hepatotoxic effects. The method can be subdivided into three stages. At the first stage, a germination activation method is introduced which includes soaking the spores in a solution to induce germination, and placing the germination treated spores in a culture box to induce the synthesis of bioactive substances and softening of the cell walls of the spores. At the second stage, sporoderm-broken ganoderma spores are collected by treating the episporos with cell wall breaking enzymes and/or mechanical force. At the last stage, the bioactive substances are extracted from the sporoderm-broken spores by drying at low temperature followed by extraction.

10 Claims, No Drawings